

12/17/99



5/17/01
207597/60
OL SR 81537

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. 2047.114

First Named Inventor or Application Identifier

SUBBARAO V. PONAKALA ET AL.

Express Mail Label No.

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

☐ Fee Transmittal Form
(Submit an original, and a duplicate for fee processing)

2. ☒ Specification Total Pages 63

3. ☒ Drawing(s) (35 USC 113) Total Sheets 4

4. ☒ Oath or Declaration Total Pages 2

a. ☐ Newly executed (original or copy)

b. ☒ Unexecuted for information purposes

c. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)

[Note Box 5 below]

i. ☐ **DELETION OF INVENTOR(S)**

Signed Statement attached deleting inventor(s)
named in the prior application, see 37 CFR
1.63(d)(2) and 1.33(b)

5. ☐ Incorporation By Reference (useable if Box 4c is checked)
The entire disclosure of the prior application, from which a copy of the
oath or declaration is supplied under Box 4c, is considered as being
part of the disclosure of the accompanying application and is hereby
incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)

7. Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)

a. ☐ Computer Readable Copy

b. ☐ Paper Copy (identical to computer copy)

c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))

9. ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)

10. ☐ English Translation Document (if applicable)

11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS
Citations

12. ☐ Preliminary Amendment

13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)

14. ☐ Small Entity Statement(s) ☐ Statement filed in prior application
Status still proper and desired

15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)

16. ☐ Other: _____

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No. /

18. CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label

05514
(Insert Customer No. or Attach bar code label here)

or ☐ Correspondence address below

NAME

Address

City

State

Zip Code

Country

Telephone

Fax

CLAIMS	(1) FOR	(2) NUMBER FILED	(3) NUMBER EXTRA	(4) RATE	(5) CALCULATIONS
	TOTAL CLAIMS (37 CFR 1.16(c))	53-20 =	33	X \$ 18.00 =	\$ 594.00
	INDEPENDENT CLAIMS (37 cfr 1.16(b))	2-3 =	0	X \$ 78.00 =	\$ 0
	MULTIPLE DEPENDENT CLAIMS (if applicable) (37 CFR 1.16(d))			\$260.00 =	\$ 0
				BASIC FEE (37 CFR 1.16(a))	\$760.00
			Total of above Calculations =		\$1354.00
	Reduction by 50% for filing by small entity (Note 37 CFR 1.9, 1.27, 1.28).				
	TOTAL =				\$1354.00

19. Small entity status

- a. ☐ A Small entity statement is enclosed
- b. ☐ A small entity statement was filed in the prior nonprovisional application and such status is still proper and desired.
- c. ☐ Is no longer claimed.

20. ☒ A check in the amount of \$ 1354.00 to cover the filing fee is enclosed.21. ☐ A check in the amount of \$ _____ to cover the recordal fee is enclosed.

22. The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 06-1205:


- a. ☒ Fees required under 37 CFR 1.16.
- b. ☒ Fees required under 37 CFR 1.17.
- c. ☐ Fees required under 37 CFR 1.18.

SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED

NAME

RAYMOND R. MANDRA (Reg. No. 34,382)

SIGNATURE



DATE

December 16, 1999

- 1 -

TITLE

N-[N-(3,3-DIMETHYLBUTYL)-L- α -ASPARTYL]-L-PHENYLALANINE
1-METHYL ESTER AS A SWEETENER IN CHEWING GUM

5

This application claims the benefit of U.S. Provisional
Patent Application No. 60/112,915, filed December 18,
1998.

10

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates to the use of N-[N-(3,3-
15 dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl
ester (neotame) as a sweetener in chewing gum. The use
of neotame as a sweetener in chewing gum results in
prolonged sweetness and flavor and low concentration of
sweetener, as compared to the use of other high-
20 intensity sweeteners.

Related Background Art

Chewing gums comprise many different ingredients including both natural and high-intensity sweeteners.

- 5 Advantages of using high-intensity sweeteners in chewing gum include flexibility of sweetness taste delivery and economics. Among high-intensity sweeteners, aspartame has been the most widely used in chewing gum formulations. U.S. Patent Nos. 3,943,258,
10 3,982,023, 4,036,992, 4,064,274, 4,122,195, 4,139,639, 4,158,068, 4,246,286, 4,248,894, 4,252,830, 4,317,837, 4,357,354, 4,374,858, 4,382,963, 4,384,004, 4,556,565, 4,612,195, 4,673,577, 4,704,288, 4,711,784, 4,722,844, 4,722,845, 4,738,854, 4,741,910, 4,774,094,
15 4,800,095, 4,816,265, 4,822,621, 4,822,622, 4,828,845, 4,929,447, 5,064,658, 5,110,608, 5,126,151, 5,139,797, 5,167,972, 5,175,009, 5,192,561, 5,221,543, 5,296,244, 5,334,397, 5,399,365, 5,425,961, 5,431,929, 5,458,892, 5,466,471, 5,470,566, 5,510,123, 5,536,510, 5,612,070,
20 WO 96/20608, WO 98/03076, EP 196640, JP 60-102147, JP 58-198250, JP 58-19034, and EP 27024 all relate to the use of aspartame in chewing gum. Other high-intensity sweeteners that have been used include acesulfame-K, cyclamates, saccharin, and sucralose. Blends of these
25 sweeteners have also been used in chewing gums in attempts to achieve overall improvements of sweetness properties.

- One problematic aspect of using high-intensity
30 sweeteners in chewing gum is the rapid release of the sweetener relative to the length of time consumers prefer to chew the gum. More specifically, each high-

intensity sweetener is associated with its own inherent sweetness release characteristic. It is possible to compensate for this by using high levels of sweeteners; in this way, sweetness is perceived by the consumer to last longer because more sweetener is present while chewing. However, the use of sweeteners in this fashion may result in a chewing gum which has too high of an initial sweetening impact or may result in adverse taste effects such as bitterness or other off-
10 notes. The use of high levels of high-intensity sweeteners may also require processing and formulation changes which relate to higher cost concerns. Thus, the ability to extend the sweetness of a chewing gum composition without compromising sweetness delivery and
15 economics is still a sought-after goal of chewing gum manufacturers.

There have been many attempts to extend the sweetness delivery in chewing gum compositions using controlled
20 release methods. In most cases, the sweetener is "bound" to other ingredients and are then released to the palate in a more controlled fashion while chewing. For example, WO 96/20608 describes the use of a multiple encapsulation method wherein aspartame and
25 acesulfame-K are first entrapped on a coating by absorption or agglomeration. U.S. Patent 4,997,659 teaches alitame encapsulation processes for the purpose of delayed release. WO 99/13870 describes a method for producing chewing gum containing acyclic carboxamides
30 applicable to sweeteners and flavors in order to produce a modified release when the gum is chewed. Although these methods provide improvement in the

extension of sweetness while the gum is chewed, the controlled release results from either the addition of encapsulated ingredients or modification of gum components. A chewing gum composition where no
5 encapsulation or modification is required for extending the sweetness would be highly desirable to both the consumer and the chewing gum manufacturer.

Another aspect which consumers find undesirable in
10 chewing gum compositions is the rapid release of flavor while the gum is being chewed. Generally, flavors are released from the chewing gum at a slower rate compared to that of the sweeteners due to the difference in the flavor's solubility between the gum base and saliva.
15 Many attempts to extend the flavor of chewing gum have been made using encapsulation, agglomeration, and/or absorption methods. For example, U.S. Patent No. 5,217,735 describes using modified cellulose to absorb flavor ingredients and then incorporating the cellulose
20 into chewing gum. U.S. Patent No. 5,156,866 describes flavor component particles coated with sterol. The use of particles and resins where flavor ingredients are either agglomerated or absorbed are taught in U.S. Patent No. 5,116,627, EP 320522 and WO 91/01132.
25 Though these methods offer some improvement, increased flavor extension without the use of modifying methods is still desired by chewing gum manufacturers and the consumer.

30 Yet another problem with chewing gum compositions which comprise dipeptide sweeteners is shelf-life stability. Attempts to increase the shelf-life stability of

aspartame and alitame in chewing gum compositions have been made since the introduction of these sweeteners. Presumably interactions between aspartame and alitame with other chewing gum components take place over time, rendering both aspartame and alitame susceptible to degradation. A number of references describe materials and methods of encapsulation for this purpose. For example, WO 96/20608 describes an encapsulation method to enhance the stability of aspartame in chewing gum when used in a sweetener blend with acesulfame-K; WO 95/17828 teaches the use of beta-glucans; U.S. Patent No. 5,277,919 describes the use of benzaldehyde acetals; and U.S. Patent No. 4,822,621 describes the use of organic acids. In each case, protection methods are described which provide improvement of the shelf-life stability of either aspartame or alitame. While some improvement of shelf-life stability has resulted, it is still desirable to use various encapsulation materials and methods in ways which offer even greater shelf-life stability.

SUMMARY OF THE INVENTION

This invention is directed to chewing gum compositions sweetened with neotame.

The invention also relates to methods of sweetening a chewing gum composition with neotame.

In a preferred embodiment of the present invention, the neotame is encapsulated before incorporation in a chewing gum composition.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a plot of the sweetness intensity of chewing gum prepared with sugar (sucrose), saccharin, aspartame, acesulfame-K, sucralose or neotame as a function of chewing time. Each chewing gum comprises the corresponding sweetener at a sweetener concentration that provides an identical ("isosweet") degree of sweetness. The sweetness intensity was measured by a trained panel of experts and is expressed as the amount of sweetness based on a sweetness intensity scale of 0-15 units.

FIG. 2 is a plot of peppermint flavor intensity verses chewing time of chewing gums prepared with isosweet levels of aspartame (3000 ppm) and neotame (100 ppm). The chewing gum prepared with aspartame comprised a 1.5 wt% peppermint flavor, while the four neotame chewing gums comprised peppermint flavor at concentrations of 1.5, 1.25, 1.00, and 0.75 wt%. The peppermint flavor intensity was measured by a trained panel of experts and is plotted as a function of chewing time. The peppermint flavor intensity is expressed as the amount of peppermint flavor based on a universal peppermint intensity scale of 0-15 units.


FIG. 3 is a plot of both sweetness and flavor intensity of chewing gums prepared with isosweet amounts of aspartame (3000 ppm) and neotame (100 ppm), each comprising 1.5% peppermint flavor. The sweetness and peppermint flavor were measured by a trained panel of experts and are expressed as the amount of sweetness

and peppermint flavor based on corresponding intensity scales of 0-15 units.

FIG. 4 is a plot of sweetness intensity verses chewing
 5 time of chewing gums prepared with neotame and
 comprising different flavor concentrations of 1.5,
 1.25, 1.00, and 0.75 wt%. The sweetness intensity was
 measured by a trained panel of experts and is plotted
 as a function of chewing time. The sweetness intensity
 10 is expressed as the amount of sweetness based on a
 universal sweetness intensity scale of 0-15 units.

DETAILED DESCRIPTION

15 According to the present invention, N-[N-(3,3-
 dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl
 ester (neotame), a high potency dipeptide sweetener
 (about 8000x sweeter than sucrose) that has the formula



is used as a sweetener in chewing gum. It has been
 found that neotame delivers an acceptable level of

sweetness in chewing gum and, in fact, prolongs
sweetness and flavor, compared to any other sweetener.
Additionally, because neotame is a high potency
sweetener, it may be used at extremely low levels
5 without sacrificing effectiveness.

Thus, one embodiment of the present invention is
directed to a sweetened chewing gum composition,
containing an effective amount of neotame to sweeten
10 the chewing gum.

The sweetness profile of neotame in chewing gum is
illustrated in FIG. 1. Also illustrated in FIG. 1 is a
comparison of the sweetness intensity profiles, as
15 measured by a trained panel of experts, of chewing gums
prepared with neotame, aspartame, acesulfame-K,
saccharin, sucralose, and sugar (sucrose). The
sweetness detected is expressed as the amount of
sweetness based on a universal sweetness scale of 0-15,
20 and the values were obtained by having subjects of the
trained panel chew pieces of gum containing the
sweetener and reporting perceived sweetness after
various chewing intervals. Since each chewing gum was
prepared with an "isosweet" concentration (i.e., same
25 level of sweetness; e.g., 100 ppm neotame and 3000 ppm
aspartame are isosweet concentrations) of the
respective sweetener and the scale is a measure of
absolute sweetness at any point during chewing, the
different sweetener profiles can be compared directly
30 to each other.

The evaluation shows that the initial rate of sweetness detection that takes place between the start of chewing and 2 minutes is greater for the chewing gum comprising sucrose than the chewing gums comprising the other
5 sweeteners. The maximum intensity perceived for a chewing gum comprising sucrose is around 1 minute's time, whereas all of the high-intensity sweeteners have a maximum sweetness intensity at around two minutes' time, all nearly identical to each other, but clearly
10 distinct from the sweetness intensity maximum of sucrose.

Further evaluation of the plot clearly demonstrates the sweetness extension provided by neotame, in particular,
15 that which occurs after around six minutes of chewing time, where neotame has been found to have a flatter or more uniform sweetness intensity throughout the remaining chewing time of 20 minutes. The sweetness extension provided by neotame relative to the other
20 sweeteners can also be illustrated by comparing the actual amount of sweetness intensity loss which occurs over a specific time period of chewing.

Table 1 summarizes the sweetness intensity loss rates
25 and average sweetness intensities which occur at a chewing time period of between 6 and 10 minutes and those which occur at a chewing time period between 10 and 20 minutes for a variety of sweeteners. The sweetness intensity loss rate is calculated as the
30 difference in sweetness intensities measured at any two time points divided by the number of minutes of the

time interval, and approximates the mathematical slope of the downward trend of sweetness intensity over time.

5 Table 1. Average sweetness loss rates and average sweetness intensities.

sweetener	average sweetness loss rate at 6-10 minutes (intensity units/min)	average sweetness intensity at 6-10 minutes (intensity units)	average sweetness loss rate at 10-20 minutes (intensity units/min)	average sweetness intensity at 10-20 minutes (intensity units)
neotame (100 ppm)	0.25	4.7	0.09	3.7
aspartame (3000 ppm)	0.52	4.6	0.25	2.3
acesulfame-K (3200 ppm)	0.45	4.3	0.26	2.0
saccharin (1000 ppm)	0.52	3.1	0.15	1.2
sucralose (600 ppm)	0.52	3.3	0.16	1.3
sucrose (60%)	0.45	2.1	0.11	0.6

20 For example, between 6 and 10 minutes of chewing, the sweetness intensity loss rate of neotame is determined to be 0.25 sweetness intensity units per minute (i.e. $(5.2-4.2 \text{ sweetness intensity units})/4 \text{ minutes} = 0.25$). A lower sweetness loss rate corresponds directly to a

25 greater sweetness extension. The results shown in Table 1 indicate that the sweetness loss rate of neotame is nearly half that of the sweetness loss rate of the other sweeteners evaluated within the time period of between 6 and 10 minutes. Similarly, for the

30 time period of chewing between 10 and 20 minutes, the sweetness intensity loss rate of neotame is 0.09 sweetness intensity units per minute, where again the

value for neotame is significantly less than that of the other sweeteners evaluated.

While Table 1 describes the average sweetness loss rate for a chewing gum composition having 100 ppm neotame, it is expected that similar average sweetness loss rates attributed to neotame would be exhibited in all chewing gum compositions containing neotame, regardless of the actual concentration of neotame.

10 It should be noted that the sweetness intensity provided by sucrose falls off so rapidly once the sweetness intensity maximum is reached (around 1 minute) that very little sweetness intensity is
15 detected by the panel at the later chewing times. For this reason, it is important to realize that the rate of sweetness loss is only one aspect of demonstrating the sweetness extension advantage of neotame, as the rate of sweetness intensity loss should be viewed
20 relative to the actual sweetness intensity achieved within a specific time period of chewing.

Thus, Table 1 also shows the average sweetness intensity achieved within the said time periods for
25 each chewing gum. These values were determined by calculating the average sweetness intensity measured at the specified evaluation times. For example, for the chewing gum comprising neotame at between 6 and 10 minutes, the average sweetness intensity is determined
30 to be $(5.2+4.6+4.2)/3 = 4.7$.

Comparison of the average sweetness intensities presented in Table 1 shows that neotame maintains a higher sweetness intensity while undergoing sweetness loss during the specified chewing time period. The
5 comparison of the different chewing gums clearly shows that neotame provides sweetness extension by its advantageous slower rate of sweetness loss and by its ability to maintain a higher sweetness intensity throughout the duration of chewing.

10 It should be noted that Table 1 describes the average sweetness intensity for a chewing gum composition having 100 ppm neotame. It is expected that a proportionally higher or lower sweetness intensity
15 would be maintained in chewing gums having proportionally higher or lower concentrations of neotame.

From this evaluation, the sweetness extension provided
20 by neotame is such that between the time duration of about 6 to about 10 minutes of chewing, the average sweetness loss rate is less than about 0.3 sweetness intensity units per minute and an average sweetness intensity of about 4 sweetness intensity units or
25 greater is maintained. Furthermore, at between the time duration of about 10 to about 20 minutes of chewing, the average sweetness loss rate is less than about 0.15 sweetness intensity units per minute and an average sweetness intensity of about 3 sweetness
30 intensity units or greater is maintained.

The advantage of sweetness extension provided by neotame can also be illustrated by comparison of the sweetness intensity half-lives over time. Table 2 summarizes the half-lives of sweetness intensity of chewing gums comprising different sweeteners after the maximum sweetness intensities (around two minutes) are reached. The half-life is a measure of the time of chewing in which half of the sweetness is lost relative to the time in which the sweetness intensity maximum is achieved.

Table 2. Half-lives of sweetness intensity.

sweetener	half-life (minutes)
neotame (100 ppm)	14.9
aspartame (3000 ppm)	6.4
acesulfame-K (3200 ppm)	5.5
saccharin (1000 ppm)	4.5
sucralose (600 ppm)	4.5
sucrose (60%)	3.5

For example, the chewing time in which neotame lost half of its sweetness was 14.9 minutes. The chewing time in which half of the sweetness was lost in chewing gums comprising other sweeteners was significantly less and not greater than 6.4 minutes.

The amount of sweetening ingredients including neotame in the appropriate proportions can be readily formulated to provide a desired sweetness level, as well as a desired amount of sweetness extension. For example, at a sweetness intensity less than about 2 intensity units, it is believed that sweetness is not perceived by most consumers of chewing gum. Preferably, sweetness is best perceived above a sweetness intensity of about 2-3. However, those skilled in the art will understand that the degree of sweetness perception varies greatly depending upon the individual consumer.

As previously mentioned, FIG. 1 shows that chewing gums comprising each of the high-intensity sweeteners have the same sweetness intensity maximum at around two minutes, followed by the characteristic loss of sweetness intensity over time. Chewing gum comprising sucrose, however, has a sweetness intensity maximum at around one minute, followed by a sweetness intensity loss rate similar to high-intensity sweeteners other than neotame. Due to neotame's sweetness extension properties, it is now possible to incorporate a blend of neotame and a sweetener having a more rapid sweetness impact in the chewing gum composition to provide a sweetness which has both an immediate sweet impact and extended sweetness. The use of neotame in this manner offers greater flexibility and provides a method of optimizing the overall sweet taste profile of chewing gum compositions.

100-50000-1-1700

Rapid release sweeteners contemplated by the present invention include, but are not limited to, sucrose, mannitol, high fructose corn syrup, sorbitol, dextrose, corn syrup solids, hydrogenated starch hydrolysates, 5 invert sugar (both liquid and dried forms), fructose, xylitol, and combinations thereof. Rapid release sweeteners that have a sweetness intensity maximum in a chewing gum composition within the first two minutes of chewing are preferred. Those skilled in the art will 10 understand that these sweeteners may also be present in the chewing gum to provide bulk and other functionality.

According to the present invention, a chewing gum can 15 now be prepared to provide not only an immediate and extended sweetness, but a better balance of both sweetness and flavor over an extended chewing period, a type of chewing gum heretofore unknown.

20 In that regard, neotame possesses exceptional flavor extension characteristics in chewing gum. FIG. 2 illustrates these flavor extension properties where the peppermint flavor intensities of chewing gums prepared with isosweet amounts of aspartame and neotame are 25 plotted as a function of chewing time, as measured by a trained panel of experts.

According to FIG. 2, the rate of flavor loss of the chewing gum comprising neotame having 1.5% peppermint 30 flavor is significantly less than that of the chewing gum comprising aspartame having the same amount of flavor, measured after the flavor intensity maximum has

been reached (around 4 minutes). As in the case of sweetness intensity comparisons, since the chewing gums were prepared with an identical amount of flavor and the scale is a measure of absolute flavor at any point during chewing, the different flavor profiles can be compared directly to each other. The rate of flavor intensity loss of the chewing gum comprising neotame is 0.0875 flavor intensity units per minute, while that of chewing gum comprising aspartame was determined to be 0.150 flavor intensity units per minute. Thus, the rate of flavor loss of a chewing gum comprising neotame is nearly half that of the flavor loss for a chewing gum comprising aspartame at an isosweet level and containing an equal concentration of peppermint flavor.

15

It was also discovered that chewing gums comprising neotame containing lesser amounts of peppermint flavor (i.e. 0.75%, 1.00%, and 1.25%) result in a flavor intensity loss over time that is non-linear. Since the rate of flavor intensity loss is not linear over the duration of the chewing time with respect to the actual flavor concentration used in the chewing gums, the chewing gums comprising neotame with lesser amounts of peppermint flavor show the advantage of providing a reduced flavor intensity rate loss while maintaining a greater flavor intensity. This advantage provided by neotame in chewing gum can clearly be seen from the summary of flavor intensity loss rates as shown in Table 3.

30

CONFIDENTIAL

Table 3. Summary of peppermint flavor loss rates at corresponding average peppermint flavor intensities for the chewing time period of between 4 and 20 minutes, as measured by a trained panel of experts.

5	sweetener	% peppermint flavor	flavor loss rate at 4-20 minutes	average flavor intensity at 4-20 minutes
	aspartame (3000 ppm)	1.50	0.150	4.2
	neotame (100 ppm)	1.50	0.088	4.6
10	neotame (100 ppm)	1.25	0.088	4.2
	neotame (100 ppm)	1.00	0.075	4.0
15	neotame (100 ppm)	0.75	0.070	3.9

Compared to that of chewing gums comprising aspartame having 1.5% flavor, the chewing gums comprising neotame at each flavor amount all have significantly reduced flavor intensity loss rates. The average flavor intensities of the chewing time period between 4 and 20 minutes are shown. For a chewing gum comprising neotame having 1.5% flavor, the flavor intensity rate loss while maintaining a greater flavor intensity is clearly advantageous compared to chewing gums comprising aspartame having the same flavor amount. Furthermore, for the chewing gums comprising neotame having less flavor amounts, the average flavor intensities are all around 4.0 intensity units, demonstrating both the flavor extension and flavor "sparing" properties of neotame.

The flavor sparing property of neotame allows for chewing gum compositions in which less flavor amounts are used to achieve a flavor intensity profile typically expected from a full-flavored chewing gum (i.e., chewing gums containing the typical or non-reduced amount of flavor). In this way, the overall cost of the chewing gum manufacture is lowered. These unique flavor extension and flavor sparing properties of chewing gums comprising neotame provide advantages in chewing gum heretofore unknown.

Based on the newly discovered flavor extension and flavor sparing properties of neotame in chewing gum, between the time duration of about 4 minutes to about 20 minutes of chewing, the flavor extension provided by neotame in a full-flavored chewing gum is such that the average flavor loss rate is less than about 0.1 flavor intensity units per minute and an average flavor intensity of about 4 flavor intensity units or greater is maintained. For chewing gums comprising neotame containing flavor amounts less than that required for a full-flavored chewing gum, the amount of flavor can be reduced from that used in a full-flavored chewing gum to at least around 50%, while providing an average flavor loss rate of less than about 0.1 flavor intensity units per minute and maintaining an average flavor intensity of about 3.5 flavor intensity units or greater.

One problematic aspect of typical chewing gums is that the intensities of both sweetness and flavor are not comparable throughout the chewing duration. The

intensity of both sweetness and flavor perceived by consumers should be both comparable and compatible at any time point of chewing. Perceived sweetness without accompaniment of flavor or perceived flavor without the
5 accompaniment of sweetness is undesirable to the consumer.

Chewing gums comprising neotame provide the advantage of maintaining a better balance between sweetness and
10 flavor compared to chewing gums comprising other sweeteners. FIG. 3 illustrates this property by comparing the sweetness and flavor profiles of chewing gum comprised of isosweet amounts of neotame and aspartame, each having identical peppermint flavor
15 amounts of 1.5%. Because of aspartame's relatively high sweetness intensity loss rate that occurs between 6 and 20 minutes compared to the loss rate of flavor intensity, the balance of sweetness intensity and flavor intensity is not maintained, as reflected by the
20 differences of sweetness and flavor intensities at chewing times greater than around 6 minutes.

However, in chewing gums comprising neotame, the rates of sweetness and flavor loss during chewing are more
25 similar, and as a result the chewing gum provides a greater balance of sweetness and flavor intensities. Based on this discovery, a chewing gum comprising neotame can be prepared wherein the difference between sweetness intensity and flavor intensity is at least
30 less than around 1 universal intensity units during the chewing time period of between around 6 minutes and around 20 minutes. The balance of sweetener extension

and flavor extension is comparably maintained throughout the duration of chewing.

A chewing gum comprising a sweetener which provides a useful level of sweetness intensity extension and allows for a significant reduction of flavor amount is also desirable. Typically, chewing gums are formulated to contain relatively large concentrations of a flavor ingredient compared to the amount of sweetness that is delivered in order to help offset the higher rate of sweetness loss while the gum is chewed. FIG. 4 illustrates the advantage provided by neotame wherein less flavor amounts are required to maintain comparable sweetness intensities and sweetness extension, in particular at extended chewing times. The measured sweetness intensities are plotted as a function of chewing time of chewing gum prepared with neotame having lesser amounts (i.e. 0.75%, 1.00%, and 1.25%) of peppermint flavor.

According to FIG. 4, the lesser amounts of peppermint flavor used do not significantly affect the sweetness intensities of neotame over the time duration of chewing, as the level of sweetness intensity of each chewing gum over the duration of the chewing time is nearly identical. Therefore, the use of neotame can provide a chewing gum composition which can contain a significantly less amount of flavor ingredient to achieve a comparable sweetness intensity that would be expected from a full-flavored chewing gum. Based on this discovery, the amount of flavor in such a gum can be reduced by at least around 50% when compared to that

of a full-flavored chewing gum sweetened with sucrose or another high-intensity sweetener.

Table 4 summarizes the half-lives of both sweetness and
5 flavor of chewing gums prepared with aspartame (3000 ppm) with 1.5 wt% flavor and neotame (100 ppm) with 1.5, 1.25, 1.00, and 0.75 wt% flavor amounts. The half-life is the time in minutes for the intensity to fall to half of its maximum value (sweetness, 2
10 minutes; flavor, 4 minutes).

Table 4. Summary of the half-lives of sweetness intensity and peppermint flavor intensity.

	sweetener	sweetness	peppermint
15	aspartame (1.50% flavor)	7.6	19.7
	neotame (1.50% flavor)	17.0	44.3
20	neotame (1.25% flavor)	17.2	31.3
	neotame (1.00% flavor)	20.7	31.7
25	neotame (0.75% flavor)	16.9	33.1

Comparing the chewing gums prepared with aspartame and neotame at the 1.5% flavor concentration clearly shows the advantage of both sweetness and flavor extension
30 provided by neotame. The sweetness extension which occurs in chewing gums containing neotame with reduced flavor amounts is also shown to be advantageous, as the half-lives are all comparable to that measured for the chewing gum containing 1.5% flavor. The flavor

extension properties that neotame provides are also shown by comparing the half-lives of flavor provided by the lesser flavor amounts to that provided by aspartame.

5

The chewing gum comprising neotame of the present invention, when chewed, releases its ingredients in a manner which yields a desirable balance of sweetness and flavor throughout the chewing period disclosed.

- 10 The chewing gum comprising neotame of the present invention has a more overall intense sweetness and flavor at longer chewing times than heretofore known compared to other sweeteners and maintains its sweetness and flavor throughout the aforementioned
- 15 chewing period.

When used in chewing gum compositions, neotame possesses unique sweetness and flavor extension characteristics, as well as advantageous flavor sparing

20 and sweetness and flavor balancing properties, most likely due to neotame's unique functionality in a given chewing gum composition. Neotame's surprising properties are likely due to favorable interactions with the gum base components. Also, since neotame

25 displays a relatively high solubility in non-aqueous solvents compared to other high-intensity and natural sweeteners, it might be expected that neotame would remain in the gum base phase, rather than quickly dissipate in the saliva phase as demonstrated by other

30 sweeteners during chewing. However, neotame does provide a significant sweetness impact during the initial chewing period in addition to maintaining a

CONFIDENTIAL

sustained sweetness intensity throughout the duration of chewing. The functionality of neotame in chewing gum is likely linked to both its solubility behavior as well as its affinity properties with the chewing gum
5 components.

The neotame suitable for use in the present invention may be prepared through a variety of methods. The chemical synthesis of neotame is disclosed in U.S.
10 Patent No. 5,480,668, U.S. Patent No. 5,510,508, and U.S. Patent No. 5,728,862, the disclosure of each of which is incorporated by reference herein. Salts and complexes of neotame, as well as blends of neotame with other high intensity and natural sweeteners, are also
15 suitable for use in the present invention.

The neotame used in chewing gum compositions may take any form. For example, it may be a salt or complex such as described in U.S. Patent Application No.
20 09/146,963, U.S. Patent Application No. 09/146,964, U.S. Patent Application No. 09/148,134, U.S. Patent Application No. 09/146,965, all filed September 4, 1998, the disclosure of each of which is incorporated by reference herein. Other exemplary forms of neotame
25 that may be used in this invention include co-crystallized forms and cyclodextrin complexes, such as described in U.S. Patent Application No. 09/154,568 and U.S. Provisional Patent Application No. 60/100,867, both filed September 17, 1998, the disclosures of each
30 of which are incorporated by reference herein. Agglomerates and other processed forms of neotame, including granulated neotame as described in U.S.

Provisional Patent Application No. 60/126,363, filed March 26, 1999, the disclosure of which is incorporated by reference herein, and various crystallized forms of neotame using different processes may also be used.

5

10 The amount of neotame can be readily formulated to provide a desired sweetness level in chewing gum compositions. Neotame is able to provide an acceptable sweetness in chewing gum when used as the sole sweetening ingredient or when used as part of a sweetener blend. If neotame is replacing all of the sweetness conventionally found in a chewing gum, the amount will range from about 10 ppm to about 1600 ppm. More preferably, the amount of neotame will range from 15 about 40 ppm to about 600 ppm. Most preferably, this range is from about 100 ppm to about 250 ppm.

20 In other words, neotame is present in a final chewing gum composition of the present invention in an amount of from about 0.001% to about 0.16% by weight, preferably in an amount of from about 0.004% to about 0.06% by weight, and most preferably in an amount of from about 0.01% to about 0.025% by weight of the composition. Of course, lesser amounts may be employed 25 when other sweeteners are present.

A further embodiment of the present invention is directed to a chewing gum comprising a gum base, an effective amount of neotame, and any other essential 30 chewing gum ingredient.

5

10

15

20

25

35

Also contemplated for use in the chewing gum compositions of the present invention is any other high-intensity sweetener. Examples of suitable high-intensity sweeteners include (A) water-soluble naturally-occurring intense sweeteners such as dihydroxychalcones, monellin, steviosides, glycyrrhizins, dihydroflavenol, and L-aminodicarboxylic acids, aminoalkenoic acid ester amides, such as those disclosed in U.S. Patent No. 4,619,834, the disclosure of which is incorporated by reference herein, and mixtures thereof; (B) water-soluble artificial sweeteners including the soluble saccharin salts of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide, the potassium salt of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide (acesulfame-K), the free acid form of saccharin, and the like, and mixtures thereof; (C) dipeptide based sweeteners including aspartame, other L-aspartic acid derivatives described in U.S. Patent No. 3,492,131, the disclosure of which is incorporated by reference herein, L-alpha-aspartyl-N-(2,2,4,4-tetramethyl-3-thietanyl)-D-alaninamide hydrate (alitame), methyl esters of L-aspartyl-L-phenyl-glycerine and L-aspartyl-L-2,5-dihydrophenyl-glycine, L-aspartyl-2,5-dihydro-L-phenylalanine, L-aspartyl-L-(1-cyclohexene)-alanine, and the like, and mixtures thereof; (D) water-soluble intense sweeteners derived from naturally occurring water-soluble sweeteners, such as chlorinated derivatives of ordinary sugar (sucrose), e.g., chlorodeoxysugar derivatives and mixtures thereof, and (E) protein based intense sweeteners such as thaumacoccus danielli (thaumatin I and II). Again, use of these sweeteners in chewing gum

compositions would naturally reduce the level of neotame required for a particular application.

These other natural and high intensity sweeteners may
5 be employed in the chewing gum compositions of the present invention as separate ingredients added at some point in the processing or as part of a blend with neotame. Those skilled in the art will recognize that any combination of natural and high intensity
10 sweeteners may be employed in the chewing gum. Further, those skilled in the art will recognize the sweetener may also function in the chewing gum in whole or in part as a water soluble bulking agent or softener. In addition, a softener may be combined with
15 the sweetener such as in an aqueous sweetener solution. In one preferred embodiment of the present invention, a blend of neotame and xylitol is used to sweeten a chewing gum composition. When used in a blend with another sweetener, the neotame is generally present in
20 an amount of from about 0.1 ppm to about 1200 ppm.

In general, a chewing gum composition comprises a water soluble bulk portion and a water insoluble chewable gum
base portion, each of which are comprised of specific
25 ingredients which make up the chewing gum composition. The water soluble portion dissipates with a portion of the sweetener and flavoring agent over a period of time during chewing. The gum base portion is retained in the mouth throughout the chew. As can be appreciated
30 by one of ordinary skill in this art, any of the materials listed at 37 C.F.R. §172.615 may be

incorporated into the chewing gum compositions of the present invention.

The insoluble gum base generally comprises resins, fats
5 and oils, waxes, elastomers, softeners and inorganic fillers. Resins generally include polyvinyl acetate and terpene resins. Fats and oils generally include animal fats such as lard and tallow, vegetable oils such as soybean and cottonseed oils, hydrogenated and
10 partially hydrogenated vegetable oils, and cocoa butter. Commonly used waxes include petroleum waxes such as paraffin, and natural waxes such as candellia, carnuba, beeswax, and polyethylene. Preferably, the waxes have a melting point between 90°F and 230°F.
15 Elastomers may include polyisobutylene, isobutylene-isoprene copolymer and styrene butadiene rubber, as well as natural latexes such as chicle.

The insoluble gum base portion typically also includes
20 a filler component such as calcium carbonate, magnesium carbonate, talc, dicalcium phosphate, or combinations thereof. The insoluble gum base also generally is comprised of a softener ingredient such as glycerol monostearate, glycerol triacetate, and combinations
25 thereof. Optional ingredients generally include antioxidants, colors, emulsifiers, and ingredients which provide hygienic functionalities.

The water soluble portion of the chewing gum
30 composition may further comprise softeners, sweeteners, flavoring agents and combinations thereof. Softeners are added to the chewing gum in order to optimize the

chewability and mouthfeel of the gum. Softeners, also known in the art as plasticizers or plasticizing agents, generally constitute between about 0.5 to about 15.0 weight percent of the chewing gum. Softeners contemplated by the present invention include glycerin, lecithin, and combinations thereof. Further, aqueous sweetener solutions such as those containing sorbitol, hydrogenated starch hydrolysates, corn syrup and combinations thereof may be used as softeners and binding agents in gum.

Texturizers suitable for use in the present invention include, without limitation, lycasin, glycerin, mannitol, and combinations thereof.

Humectants suitable for use in the present invention include, without limitation, glycerin, sorbitol, and combinations thereof.

Crystallization inhibitors suitable for use in the present invention include, without limitation, mannitol.

Bulking or binding agents suitable for use in the present invention include, without limitation, dextrose, maltodextrin, lactose, inulin, cellulose, cellulose derivatives, gelatin, xanthan, guar, pectins, locust bean, alginates, agar, carrageenans, gum acacia, tara gum, karaya gum, gellan gum, gurrcellaran, tragacanth, guar gum hydrolysate, ghatti, microcrystalline cellulose, carbomethyl-cellulose, hollocellulose, cellulose gel, polydextrose,

maltodextrose, isomalulose, polymaltose,
arabinogalactan, palatinose, starches, starch
hydrolysates, hydrogenated starch hydrolysates,
partially hydrolyzed starch, dextrins, hydrogenated
5 hexoses, fructooligosaccharides, sorbitol, xylitol,
mannitol, maltitol, galactitol, isomalt, and mixtures
thereof.

Organic acids suitable for use in the present invention
10 include, without limitation, citric acid, malic acid,
tartaric acid, and mixtures thereof.

Any suitable colorant may be used in the chewing gum
compositions of the present invention. There are no
15 limitations as to the types of colorant that can be
used in compositions containing neotame; therefore, any
colorant recognized in the art as a traditional
colorant useful for chewing gum compositions can be
used.

20 Any suitable flavor may be used in the chewing gum
compositions of the present invention. Chewing gum
compositions typically contain a variety of different
flavorings, and there is no limitation as to the types
25 of flavoring ingredients which can be used in chewing
gums sweetened with neotame. Flavoring ingredients are
typically present in chewing gums in amounts from about
0.1 wt% to about 10 wt%, and preferably from about 0.75
wt% to about 3 wt% of the gum. The flavoring
30 ingredient may comprise essential oils, synthetic
flavors, or mixtures thereof including, but not limited
to, oils derived from plants and fruits such as citrus

0046500-14700

oils, fruit essences, peppermint oil, spearmint oil, clove oil, oil of wintergreen, anise, cinnamon, and fruit flavors such as tutti frutti, strawberry, raspberry, lemon, and orange. Artificial flavoring components are also contemplated for the present invention. Those skilled in the art will recognize that natural and artificial flavoring ingredients may be combined in any blend that is deemed acceptable. All such flavors and flavor blends are contemplated by the present invention. Generally, the amounts of colorants and flavorings that are used are normally a matter of preference subject to type and desired end-strength.

Optional ingredients such as emulsifiers and pharmaceutical agents may also be added to the chewing gum, and these types of additives are also contemplated by the present invention.

In another preferred embodiment of the present invention, neotame is encapsulated or formed as an admixture with a suitable agent for use as a sweetener in chewing gum compositions. Incorporation of said encapsulated form or admixture form of neotame into chewing gum compositions extends the shelf-life stability of neotame. The encapsulated or admixture forms provide protection of neotame from chewing gum components which may interact with neotame over time. As a consequence, chewing gum compositions containing encapsulated or admixed neotame are more stable than chewing gums comprising neotame in a non-encapsulated or non-admixture form.

Suitable materials include, without limitation, cellulose and cellulose derivatives such as hydroxypropylmethyl cellulose, stearic acid, shellac, polyethylene wax 500, zein, sterine 27, alginates, 5 gelatin, starches, proteins, sugars, sugar alcohols, complex carbohydrates, gums, hydrocolloids, gellan gum, polydextrose, polywax, hydrogenated starch hydrolyzate, polyvinyl acetate, xanthan gum, carrageenan, dextrose, malic acid, maltodextrin, gum arabic and combinations 10 thereof. Materials preferable for use in the present invention include hydroxypropylmethyl cellulose, polyvinyl acetate and zein. In an encapsulated or admixed composition, neotame is generally present in an amount of 0.5-20% by weight of the encapsulated or 15 admixed composition.

Furthermore, those skilled in the art will recognize that any traditional method of encapsulation or that used to form an admixture can be employed. These 20 include, but are not limited to, methods such as spray drying, granulation, agglomeration, fluidized bed agglomeration, complex coacervation, spray chilling, prilling, extrusion, methods of drying such as drum-drying, freeze-drying, oven-drying, processing through 25 a rotary disk, and other similar techniques.

One skilled in the art would appreciate that any encapsulation method or material useful in the encapsulation or admixing of aspartame or other high- 30 intensity sweeteners, as well as other chewing gum ingredients, would be suitable for use in the encapsulation or admixture of neotame. Several of

these methods and materials can be found in U.S. Patent Nos. 4,139,639, 4,384,004, 4,590,075, 4,673,577, 4,711,784, 4,722,845, 4,726,953, 4,816,265, 4,824,681, 4,911,934, 4,929,447, 4,933,190, 4,952,402, 4,975,270, 5 4,978,537, 4,981,698, 4,986,991, 5,043,169, 5,064,658, 5,108,763, 5,126,151, 5,139,798, 5,154,939, 5,164,210, 5,167,972, 5,169,697, 5,192,561, 5,334,397, 5,338,809 and 5,532,004, WO 8911212, WO 8903170, WO 9007859, WO 9007864, WO 9103147, WO 960608, WO 9704662, WO 9803076, 10 EP 252374, EP 427541, and GB 2225923, the disclosures of which are incorporated by reference herein.

Other ingredients may be added to provide additional functionalities which may be desired for certain 15 applications in the same manner used with aspartame and other high-intensity sweeteners. For example, U.S. Patent Nos. 5,217,735, 5,221,543, 4,800,095, 5,112,625, 4,885,175, 4,839,184, 4,374,858, and 5,665,406 relate to the manipulation of desired functionalities of 20 aspartame and are incorporated by reference herein.

As mentioned above, it is possible to use neotame in combination with other natural or high-intensity sweeteners. In such embodiments of the present 25 invention, neotame may be encapsulated or admixed, while the other sweetener(s) is not. Alternatively, either both or neither neotame and the other sweetener(s) may be encapsulated or admixed for use in chewing gum formulations.

30

Prior to encapsulation, neotame may be spray-dried in combination with any suitable vehicle. These vehicles

include, without limitation, gum arabic, starch, and maltodextrin. Typically, these vehicles are used in approximately 20% concentration in water and neotame is generally present in an amount of 2-10% by weight of the spray-dried composition.

Those skilled in the art will recognize that various materials and methods of encapsulation will afford different degrees of protection, thereby resulting in different levels of shelf-life stability enhancement of neotame. The preferred quantity and type of encapsulating or other material employed to protect neotame from interacting with other chewing gum components may be specifically tailored depending upon a variety of factors including, but not limited to, the encapsulation or other method employed to form said admixture, the type of encapsulant or material agent used to form said admixture, and the amount of encapsulant or other material agent used to form said admixture. Since the level of protection may be governed by a number of factors such as cost, desired stability enhancement, and degree of processing conditions, it is contemplated that those skilled in the art will recognize the various factors and make adjustments as necessary in order to achieve a level of stability enhancement desired.

As part of the present invention, it is contemplated that the sweetness extension properties of neotame during the chewing period can be increased further by adding said neotame to chewing gum in a form that affords an even longer sweetness extension, such as

that expected from encapsulated, co-dried, and agglomerated forms.

It is also contemplated by the present invention that
5 chewing gum comprising neotame can be prepared by these
methods to yield various controlled rates of sweetness
intensity over the duration of a particular chewing
time. Thus, manufacturers will be able to produce a
chewing gum which may more closely approximate a
10 constant desirable sweetness level over time. Examples
of extending the sweetness in chewing gums by
encapsulation or forming an admixture with another
substance are well known in the art, and it is
contemplated that this type of formulation can be
15 carried out by one ordinarily skilled in the art.

It is also contemplated by the present invention that
one ordinarily skilled in the art will recognize that
various forms of chewing gums can be manufactured. In
20 general, chewing gum is manufactured by sequentially
adding the various chewing gum ingredients to any
commercially available mixer known in the art. After
the ingredients have been thoroughly mixed, the gum
mass is discharged from the mixer and shaped into the
25 desired form such as by rolling into sheets and cutting
into sticks, extruding into chunks, or casting into
pellets.

It is also anticipated that the physical form in which
30 neotame exists when added to the chewing gum
composition may affect its sweetness extension
properties during chewing. Different physical forms

contemplated include, but are not limited to, powdered, granular, extruded, compacted, or particulate forms. Chemical forms contemplated include anhydrous forms, amorphous forms, partially hydrated forms, and solvated forms. Of course, it is contemplated that the sweetness extension of these forms can further be altered by encapsulation or agglomeration methods, or co-drying to provide an admixture of neotame and another acceptable chewing gum component.

It is also contemplated that neotame may be altered in its sweetness and flavor extension properties by dissolving said neotame in a food acceptable solvent prior to being added to the chewing gum composition during formulation. Suitable solvents include, but are not limited to, ethanol, propylene glycol, glycerin, glycerol triacetate, vegetable oil, ethyl acetate, and combinations thereof. In addition, flavoring agents, because of their ability to dissolve neotame, individually or as a blend, are considered suitable solvents. The optimal solubility level of neotame in a solvent that is amenable to the manufacture of chewing gum can be readily determined.

Another embodiment of the present invention is directed to a process for sweetening a chewing gum composition by including in the chewing gum composition an effective amount of neotame. One of ordinary skill in this art would appreciate that the inclusion of neotame in a chewing gum composition can be accomplished by any means. Suitable means include adding neotame to a mixture of chewing gum ingredients, adding neotame to

the surface of a chewing gum composition and then adding a protectant such as zein or a polyol, and adding a liquid dispersion of neotame in a substance such as glycerin to a piece of chewing gum.

5

All of the aforementioned advantages are made possible by the present invention disclosed herein. Other advantages realized from the present invention which are not mentioned but which will be readily apparent to those skilled in the art are also contemplated.

10

The art has not investigated sweetener extension of neotame as a function of gum base composition. The various components of the chewing gum composition are believed to have no deleterious effect on the sweetness extension properties of neotame.

15

It is expected that a wide range of changes and modifications to the preferred embodiments described above will be apparent to those skilled in the art. The examples which follow are intended as an illustration of certain preferred embodiments of the invention, and no limitation of the invention is implied.

25

EXAMPLE 1

Chewing Gum Composition Sweetened With 0.025% Neotame

30

A chewing gum containing neotame was prepared in accordance with the present invention. The chewing gum was prepared by mixing powdered neotame with the other

ingredients listed and in the proportions specified below by the following general procedure:

	ingredient	weight percent
5	sorbitol	46.750
	gum base	26.250
	lycasin	14.583
	glycerin	5.833
	mannitol	5.100
10	peppermint flavor	1.458
	neotame	0.025

An Aaron gum mixer steam jacket was first warmed to 120-140°F. The gum base was then added to the mixer and softened. Half of the sorbitol powder was then added, followed by mixing for about two minutes. The lycasin was then added, followed by additional mixing for two minutes. Glycerin was then added, and the mixture was mixed for three minutes. The remainder of the sorbitol powder and the neotame were then added, followed by mixing for two minutes. Mannitol was then added, mixed for five minutes, followed by addition of the peppermint flavor. After mixing an additional four minutes, the gum was removed from the mixer and dusted with mannitol. The gum was then rolled and pressed to form a flat sheet of approximately 0.2 inches thick, after which the gum was cut into pieces that measured approximately 1.5 x 0.5 inches and weighed 2.6 to 2.8 grams. The pieces of gum were individually wrapped in a laminate consisting of foil, adhesive, and tissue paper. The resulting chewing gum was tested by a panel

of experts and the chewing gum exhibited both sweetness and flavor extension properties in accordance with the present invention.

5 EXAMPLE 2

Chewing Gum Composition Sweetened With 0.010% Neotame

A chewing gum containing neotame was prepared in accordance with the procedure of Example 1 using the ingredients listed below.

	ingredient	weight percent
	sorbitol	46.765
	gum base	26.250
	lycasin	14.583
15	glycerin	5.833
	mannitol	5.100
	peppermint flavor	1.458
	neotame	0.010

20 The resulting chewing gum was tested by a panel of experts, and the chewing gum exhibited both sweetness and flavor extension properties in accordance with the present invention.

25 EXAMPLE 3

Formation Of Gum Arabic, Starch, And Maltodextrin
Admixtures Of Neotame

Neotame was formed as an admixture with gum arabic,
30 starch, and maltodextrin in accordance with the present invention by mixing neotame separately with the co-

agents followed by spray-drying. Gum arabic, starch, and maltodextrin were separately prepared as 20% solutions in water. Neotame was first dissolved in ethanol, and the ethanol solution containing neotame was added to the co-agent solution. The ethanol solution containing neotame and the co-agent solution were mixed together, and the solution was then spray-dried at an outlet temperature of 130-135°F to produce a white to off-white free-flowing powder. Chewing gums comprising each admixture form were prepared in accordance with the procedure of Example 1. The resulting chewing gums were tested by a panel of experts and the chewing gums exhibited both sweetness and flavor extension properties in accordance with the present invention.

EXAMPLE 4

Agglomeration of Neotame with Dextrose

A neotame solution was prepared by dissolving 30 g of neotame into 100 g of water and 40 g of ethanol. The neotame solution was sprayed onto 970 g of dextrose in a fluid bed at a rate of 5 g/min, until all the solution had entered the fluid bed and was agglomerated to the dextrose. The resultant product was neotame diluted with dextrose containing 2.8 wt/wt% neotame. The particle size distribution for the product was: 1.2% 40 mesh, 5.9% 100 mesh, 24.5% 140 mesh, 58.4% 200 mesh and 10.0% greater than 200 mesh.

EXAMPLE 5

Encapsulation of Neotame with Polyvinyl Acetate

- 5 A 15% solution of polyvinyl acetate in methanol was prepared as the spray solution for a Wurster column. Neotame (500.0 g) diluted with dextrose from Example 4 was added to the bowl of the Wurster column. When the contents of the bowl reached a temperature of 140°F, the 15% polyvinyl acetate solution was applied to the contents of the bowl. The spray rate was maintained around 5.0 g/min, until 20% polyvinyl acetate solids was added to the contents in the bowl. The product obtained was a coating of 20% polyvinyl acetate solids on the neotame diluted with dextrose (1.83% neotame).
10 A sample of 200.0 g was collected.

- Then 300.0 g of the 20% polyvinyl acetate solids product was returned to the bowl and coated in the same manner above until a 40% polyvinyl acetate solids
20 product was obtained. A sample of 200.0 g was collected. The particle size distribution for the product was: 3.3% 40 mesh, 72.4% 100 mesh, 18.9% 140 mesh, 5.4% 200 mesh and 0.5% greater than 200 mesh.

25 EXAMPLE 6

Encapsulation of Neotame with Hydroxypropylmethyl Cellulose

- A 10% solution of hydroxypropylmethyl cellulose in
30 water was prepared as the spray solution for a Wurster column. Neotame (500.0 g) diluted with dextrose from Example 4 was added to the bowl of the Wurster column.

When the contents of the bowl reached a temperature of 170°F, the 10% hydroxypropylmethyl cellulose solution was applied to the contents of the bowl. The spray rate was maintained around 5.0 g/min, until 10% hydroxypropylmethyl cellulose solids was added to the contents in the bowl. The product obtained was a coating of 10% hydroxypropylmethyl cellulose solids on the neotame diluted with dextrose. A sample of 200.0 g was collected.

10 Then 273.0 g of the 10% hydroxypropylmethyl cellulose solids product was returned to the bowl and coated in the same manner above until a 20% hydroxypropylmethyl cellulose solids product was obtained (2.14% neotame).
15 A sample of 200.0 g was collected. The particle size distribution for the product was: 18.7% 40 mesh, 55.4% 100 mesh, 8.8% 140 mesh, 8.1% 200 mesh and 9.7% greater than 200 mesh.

20 EXAMPLE 7

Encapsulation of Neotame with Zein

A 13.5% solution of zein in methanol/water (69% methanol/16% water/13.5% zein/1.5% myvacet) was prepared as the spray solution for a Wurster column. Neotame (500.0 g) diluted with dextrose from Example 4 was added to the bowl of the Wurster column. When the contents of the bowl reached a temperature of 165°F, the 13.5% zein solution was applied to the contents of the bowl. The spray rate was maintained around 5.0 g/min, until 10% zein solids was added to the contents in the bowl. The product obtained was a coating of 10%

zein solids on the neotame diluted with dextrose. A sample of 200.0 g was collected.

- Then 273.0 g of the 10% zein solids product was
- 5 returned to the bowl and coated in the same manner above until a 20% zein solids product was obtained (2.33% neotame). A sample of 200.0 g was collected. The particle size distribution for the product was: 0.5% 40 mesh, 2.3% 100 mesh, 13.7% 140 mesh, 47.8% 200
- 10 mesh and 35.9% greater than 200 mesh.

EXAMPLE 8

Encapsulation of Neotame with Capsul Starch

- 15 Capsul starch (3 kg) (National Starch & Chemical, Bridgewater, NJ) was mixed into 7 kg water. The mixture was heated to 190°F and held for one hour. After one hour, the mixture was cooled to 150°F. Then 36.1 g neotame was added to 4.02 kg starch solution.
- 20 The resultant mixture was spray dried using a spray drier. The inlet air temperature ranged from 160°C-330°C, and the outlet temperature ranged from 90°C-120°C. The final product was neotame encapsulated with capsul starch (2.64 wt/wt% neotame). The particle size
- 25 distribution for the product was: 0.1% 40 mesh, 0.4% 100 mesh, 0.5% 140 mesh and 99.0% greater than 200 mesh.

EXAMPLE 9

Encapsulation of Neotame with Capsul Starch/Zein

5 A 13.5% solution of zein in methanol/water (69%
methanol/16% water/13.5% zein/1.5% myvacet) was
prepared as the spray solution for a Wurster column.
Neotame (400.0 g) encapsulated with capsul starch from
Example 8 was added to the bowl of the Wurster column.
When the contents of the bowl reached a temperature of
10 165°F, the 13.5% zein solution was applied to the
contents of the bowl. The spray rate was maintained
around 5.0 g/min, until 10% zein solids was added to
the contents in the bowl. The product obtained was a
coating of 10% zein solids on the neotame encapsulated
15 with capsul starch (2.42% neotame). A sample of 200.0
g was collected. The particle size distribution for
the product was: 0.6% 40 mesh, 2.1% 100 mesh, 2.6% 140
mesh, 79.8% 200 mesh and 16.0% greater than 200 mesh.

20 EXAMPLE 10

Encapsulation of Neotame with Capsul Starch/Polyvinyl
Acetate

25 A 15% solution of polyvinyl acetate in methanol was
prepared as the spray solution for a Wurster column.
Neotame (400.0 g) encapsulated with capsul starch from
Example 8 was added to the bowl of the Wurster column.
When the contents of the bowl reached a temperature of
140°F, the 15% polyvinyl acetate solution was applied
30 to the contents of the bowl. The spray rate was
maintained around 5.0 g/min, until 10% polyvinyl
acetate solids was added to the contents in the bowl.

2025 RELEASE UNDER E.O. 14176

The product obtained was a coating of 10% polyvinyl acetate solids on the neotame encapsulated with capsul starch (2.29% neotame). A sample of 200.0 g was collected. The particle size distribution for the
5 product was: 1.1% 40 mesh, 3.5% 100 mesh, 3.0% 140 mesh, 60.9% 200 mesh and 32.4% greater than 200 mesh.

EXAMPLE 11

Encapsulation of Neotame with Capsul Starch/
10 Hydroxypropylmethyl Cellulose

A 10% solution of hydroxypropylmethyl cellulose in water was prepared as the spray solution for a Wurster column. Neotame (400.0 g) encapsulated with capsul
15 starch from Example 8 was added to the bowl of the Wurster column. When the contents of the bowl reached a temperature of 170°F, the 10% hydroxypropylmethyl cellulose solution was applied to the contents of the bowl. The spray rate was maintained around 5.0 g/min,
20 until 10% hydroxypropylmethyl cellulose solids was added to the contents in the bowl. The product obtained was a coating of 10% hydroxypropylmethyl cellulose solids on the neotame encapsulated with capsul starch (2.14% neotame). A sample of 200.0 g was
25 collected. The particle size distribution for the product was: 4.6% 40 mesh, 31.7% 100 mesh, 11.3% 140 mesh, 15.6% 200 mesh and 37.8% greater than 200 mesh.

EXAMPLE 12

Encapsulation of Neotame with HiCap Starch

HiCap starch (3 kg) (National Starch & Chemical,
5 Bridgewater, NJ) was mixed into 7 kg water. The
mixture was heated to 190°F and held for one hour.
After one hour, the mixture was cooled to 150°F. Then
36.1 g neotame was added to 4.02 kg starch solution.
The resultant mixture was spray dried using a spray
10 drier. The inlet air temperature ranged from 160°C-
330°C, and the outlet temperature ranged from 90°C-
120°C. The final product was neotame encapsulated with
HiCap starch (2.73 wt/wt% neotame). The particle size
distribution for the product was: 2.5% 40 mesh, 8.0%
15 100 mesh, 2.6% 140 mesh and 87.1% greater than 200
mesh.

EXAMPLE 13

Encapsulation of Neotame with HiCap Starch/

20 Hydroxypropylmethyl Cellulose

A 10% solution of hydroxypropylmethyl cellulose in
water was prepared as the spray solution for a Wurster
column. Neotame (400.0 g) encapsulated with HiCap
25 starch from Example 12 was added to the bowl of the
Wurster column. When the contents of the bowl reached
a temperature of 170°F, the 10% hydroxypropylmethyl
cellulose solution was applied to the contents of the
bowl. The spray rate was maintained around 5.0 g/min,
30 until 10% hydroxypropylmethyl cellulose solids was
added to the contents in the bowl. The product
obtained was a coating of 10% hydroxypropylmethyl

cellulose solids on the neotame encapsulated with HiCap starch (2.30% neotame). A sample of 200.0 g was collected. The particle size distribution for the product was: 5.4% 40 mesh, 34.2% 100 mesh, 9.5% 140 mesh, 36.8% 200 mesh and 14.8% greater than 200 mesh.

EXAMPLE 14

Stability Of Encapsulated Forms Of Neotame

- 10 Chewing gum compositions using the encapsulated neotame compositions of Examples 4-12 were made in accordance with Example 2, in order to provide chewing gum samples with 100 ppm neotame. Each of the chewing gums was assayed at 4 and 8 weeks time storage time (75 °F, 55% relative humidity) for the amount of neotame present. 15 A chewing gum comprising neotame without the advantage of being encapsulated is included as a control.

Table 5. Shelf-life stability enhancement of neotame.

example (encapsulant)	% neotame remaining at 8 weeks
control	83
single coating:	
4 (dextrose)	84
8 (capsul starch)	89
12 (HiCap starch)	87
double coating:	
5 (dextrose/polyvinylacetate)	94
6 (dextrose/ hydroxypropylmethylcellulose)	100
7 (dextrose/zein)	96
9 (capsul starch/zein)	91
10 (capsul starch/ polyvinylacetate)	94
11 (capsul starch/ hydroxypropylmethylcellulose)	99

As Table 5 illustrates, the amount of neotame which remains at 4 and 8 weeks time is greater than that provided by the neotame control. Furthermore, the enhancement of stability of neotame is dependent on the type of agent used, since the range of neotame loss at the 8 week storage time varies between around 0% to around 17%. The use of hydroxypropylmethylcellulose as an encapsulation material affords the greatest stability enhancement, in which case the enhancement is around 25% measured at the 8 week storage time.

Sensory Evaluation

5 The sensory profiling of chewing gum samples was carried out using the Spectrum™ Method of Descriptive Analysis (Meilgaard, M., Civille, G.V., Carr, B.T, 1999. Sensory Evaluation Techniques, 3rd Ed. CRC Press, New York, p. 173-229). Extensive sensory training was conducted in order to familiarize the panel, consisting of 6-8 individuals, with a product category, exposing 10 the panel to the flavor attributes of the category and in-depth practice by the panelists in the use of a fifteen point scale for describing intensity differences. Once trained, the method enabled the trained panelists to indicate the sensory intensity of 15 either sweetness intensity or flavor intensity in a universal context, where 0 indicates "none" of the attribute present, 2 approximates "threshold" intensities, 5 "slight or low" level, 7.5 "moderate" intensity, 10 "strong" and 15 "extreme".

20

Chewing of the gums was initiated by the panel leader while starting a timer. As the time intervals 0.25, 0.5, 1.0, 2.0, 3.0, 4.0, 6.0, 8.0, 10.0, 12.0, 16.0, and 20.0 minutes approached, the panel leader gave 25 panelists a five second warning and then called out the time to rate the attributes. To minimize fatigue, 3-4 samples were evaluated during one session. Samples were assessed in duplicate, presented in a balanced order, unwrapped in two ounce plastic cups, and 30 randomly coded with three-digit numbers. The panelists were given a ten minute break between samples to minimize carry-over effects. Water, Mott's Natural

Applesauce and unsalted soda crackers were given as palate cleansers.

5 The data was analyzed for standard ANOVA for across sample comparison of attributes at each chew time. Data was then plotted as a graph of intensities vs. chewing time yielding either a sweetness or flavor profile for the entire chew time.

10 Other variations and modification of this invention will be obvious to those skilled in this art. This invention is not to be limited except as set forth in the following claims.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211

What is claimed is:

1. A chewing gum composition comprising N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in an amount effective to sweeten said chewing gum composition.
2. The chewing gum composition according to claim 1, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is in an amount of from about 10 to about 1600 parts per million.
3. The chewing gum composition according to claim 2, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is in an amount of from about 40 to about 600 parts per million.
4. The chewing gum composition according to claim 3, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is in an amount of from about 100 to about 250 parts per million.
5. A chewing gum composition according to claim 1, further comprising one or more additional sweeteners.
6. The chewing gum composition according to claim 5, wherein said one or more additional sweeteners is selected from the group consisting of xylose, ribulose, glucose, mannose, dextrin, galactose, fructose, sucrose, maltose, invert sugar, partially hydrolyzed starch, corn syrup solids, sorbitol, xylitol, mannitol, galactitol, maltitol, isomalt, maltodextrins,

hydrogenated starch hydrolysates, hydrogenated hexoses, hydrogenated disaccharides, dihydroxychalcones, monellin, steviosides, glycyrrhizins, dihydroflavenol, L-aminodicarboxylic acids, aminoalkenoic acid ester amides, saccharin, acesulfame, and saccharin and acesulfame salts, aspartame, alitame, chlorodeoxysugar derivatives of sucrose, thaumatin derivatives, and mixtures thereof.

7. The chewing gum composition according to claim 5, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is in an amount of from about 0.1 ppm to about 1200 parts per million.

8. The chewing gum composition according to claim 1 further comprising a chewable gum base.

9. The chewing gum composition according to claim 8, wherein said chewable gum base includes a resin which is selected from the group consisting of polyvinyl acetate and terpene resins.

10. The chewing gum composition according to claim 8, wherein said chewing gum base includes a fat or an oil which is selected from the group consisting of lard, tallow, soybean oil, cottonseed oil, hydrogenated and partially hydrogenated vegetable oils, cocoa butter, and combinations thereof.

11. The chewing gum composition according to claim 8, wherein said chewable gum base includes a wax which is selected from the group consisting of petroleum waxes,

paraffin, candellia, carnuba, beeswax, polyethylene, and combinations thereof.

12. The chewing gum composition according to claim 8, wherein said chewable gum base includes an elastomer which is selected from the group consisting of polyisobutylene, isobutylene-isoprene copolymer, styrene butadiene rubber, chicle, and combinations thereof.

13. The chewing gum composition according to claim 8 further comprising a softener which is selected from the group consisting of glycerin, lecithin, glycerol monostearate, glycerol triacetate, and combinations thereof.

14. The chewing gum composition according to claim 8, wherein said chewable gum base includes an inorganic filler which is selected from the group consisting of calcium carbonate, magnesium carbonate, talc, dicalcium phosphate, and combinations thereof.

15. The chewing gum composition according to claim 8 further comprising a texturizer which is selected from the group consisting of lycasin, glycerin, mannitol, or combinations thereof.

16. The chewing gum composition according to claim 8 further comprising a bulking or binding agent selected from the group consisting of dextrose, maltodextrin, lactose, inulin, cellulose, cellulose derivatives, gelatin, xanthan, guar, pectins, locust bean,

2025 RELEASE UNDER E.O. 14176

alginates, agar, carrageenans, gum acacia, tara gum, karaya gum, gellan gum, gurgellaran, tragacanth, guar gum hydrolysate, ghatti, microcrystalline cellulose, carbomethyl-cellulose, hollocellulose, cellulose gel, polydextrose, maltodextrose, isomalulose, polymaltose, arabinogalactan, palatinose, starches, starch hydrolysates, hydrogenated starch hydrolysates, partially hydrolyzed starch, dextrans, hydrogenated hexoses, fructooligosaccharides, sorbitol, xylitol, mannitol, maltitol, galactitol, isomalt, and mixtures thereof.

17. The chewing gum composition according to claim 8 further comprising an organic acid selected from the group consisting of citric acid, malic acid, tartaric acid, and mixtures thereof.

18. The chewing gum composition according to claim 8 further comprising a flavor selected from the group consisting of essential oils, synthetic flavors, oils derived from plants and fruits, citrus oils, fruit essences, peppermint oil, spearmint oil, clove oil, oil of wintergreen, anise, cinnamon, tutti frutti, strawberry, raspberry, lemon, orange, artificial flavorings, and combinations thereof.

19. The chewing gum composition according to claim 1, wherein between 6 and 10 minutes chewing time, the average sweetness intensity loss rate is less than 0.3 intensity units per minute.

20. The chewing gum composition according to claim 1, wherein between 10 and 20 minutes chewing time, the average sweetness intensity loss rate is less than 0.15 intensity units per minute.

21. The chewing gum composition according to claim 1, wherein a flavor is present in an amount effective to produce a full-flavored chewing gum composition and wherein between 4 and 20 minutes chewing time, the average flavor intensity loss rate is less than 0.1 intensity units per minute.

22. The chewing gum composition according to claim 1, wherein an amount of flavor used is at least 50% less than the amount of flavor in a full-flavored chewing gum composition, whereby a level of flavor intensity comparable to a full-flavored chewing gum composition is maintained.

23. The chewing gum composition according to claim 1 further comprising a rapid release sweetener.

24. The chewing gum composition according to claim 23, wherein the rapid release sweetener has a sweetness intensity maximum within the first 2 minutes of chewing.

25. The chewing gum composition according to claim 23, wherein the rapid release sweetener is selected from the group consisting of sucrose, mannitol, fructose, high fructose corn syrup, sorbitol, dextrose, corn

syrup solids, hydrogenated starch hydrolysates, invert sugar, fructose, xylitol, and combinations thereof.

26. The chewing gum composition according to claim 1, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is encapsulated by or admixed with an agent.

27. The chewing gum composition according to claim 26, wherein the agent is selected from the group consisting of cellulose, cellulose derivatives, hydroxypropylmethyl cellulose, stearic acid, shellac, polyethylene wax 500, zein, sterine 27, alginates, gelatin, starches, proteins, sugars, sugar alcohols, complex carbohydrates, gums, hydrocolloids, gellan gum, polydextrose, polywax, hydrogenated starch hydrolyzate, polyvinyl acetate, xanthan gum, carrageenan, dextrose, malic acid, maltodextrin, gum arabic and combinations thereof.

28. A process for sweetening a chewing gum composition comprising the step of:
including N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester in a chewing gum composition in an amount effective to sweeten said chewing gum composition.

29. The process of sweetening a chewing gum composition according to claim 28, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is included in an amount of from about 10 to about 1600 parts per million.

30. The process of sweetening a chewing gum composition according to claim 29, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is included in an amount of from about 40 to about 600 parts per million.

31. The process of sweetening a chewing gum composition according to claim 30, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is included in an amount of from about 100 to about 250 parts per million.

32. The process of sweetening a chewing gum composition according to claim 28 further comprising the step of including one or more additional sweeteners in said chewing gum composition.

33. The process of sweetening a chewing gum composition according to claim 32, wherein said one or more additional sweeteners is selected from the group consisting of xylose, ribulose, glucose, mannose, dextrin, galactose, fructose, sucrose, maltose, invert sugar, partially hydrolyzed starch, corn syrup solids, sorbitol, xylitol, mannitol, galactitol, maltitol, isomalt, maltodextrins, hydrogenated starch hydrolysates, hydrogenated hexoses, hydrogenated disaccharides, dihydroxychalcones, monellin, steviosides, glycyrrhizins, dihydroflavenol, L-aminodicarboxylic acids, aminoalkenoic acid ester amides, saccharin, acesulfame, and saccharin and acesulfame salts, aspartame, alitame, chlorodeoxysugar

derivatives of sucrose, thaumatin derivatives, and mixtures thereof.

34. The process of sweetening a chewing gum composition according to claim 32, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is included in an amount of from about 0.1 ppm to about 1200 parts per million.

35. The process of sweetening a chewing gum composition according to claim 28, wherein said chewing gum composition comprises a chewable gum base.

36. The process of sweetening a chewing gum composition according to claim 35, wherein said chewable gum base includes a resin which is selected from the group consisting of polyvinyl acetate and terpene resins.

37. The process of sweetening a chewing gum composition according to claim 35, wherein said chewing gum base includes a fat or an oil which is selected from the group consisting of lard, tallow, soybean oil, cottonseed oil, hydrogenated and partially hydrogenated vegetable oils, cocoa butter, and combinations thereof.

38. The process of sweetening a chewing gum composition according to claim 35, wherein said chewable gum base includes a wax which is selected from the group consisting of petroleum waxes, paraffin, candellia, carnuba, beeswax, polyethylene, and combinations thereof.

39. The process of sweetening a chewing gum composition according to claim 35, wherein said chewable gum base includes an elastomer which is selected from the group consisting of polyisobutylene, isobutylene-isoprene copolymer, styrene butadiene rubber, chicle, and combinations thereof.

40. The process of sweetening a chewing gum composition according to claim 35, wherein said chewing gum composition further comprises a softener which is selected from the group consisting of glycerin, lecithin, glycerol monostearate, glycerol triacetate, and combinations thereof.

41. The process of sweetening a chewing gum composition according to claim 35, wherein said chewable gum base includes an inorganic filler which is selected from the group consisting of calcium carbonate, magnesium carbonate, talc, dicalcium phosphate, and combinations thereof.

42. The process of sweetening a chewing gum composition according to claim 35, wherein said chewing gum composition further comprises a texturizer which is selected from the group consisting of lycasin, glycerin, mannitol, or combinations thereof.

43. The process of sweetening a chewing gum composition according to claim 35, wherein said chewing gum composition further comprises a bulking or binding agent selected from the group consisting of dextrose, maltodextrin, lactose, inulin, cellulose, cellulose

derivatives, gelatin, xanthan, guar, pectins, locust bean, alginates, agar, carrageenans, gum acacia, tara gum, karaya gum, gellan gum, gurrcellaran, tragacanth, guar gum hydrolysate, ghatti, microcrystalline cellulose, carbomethyl-cellulose, hollocellulose, cellulose gel, polydextrose, maltodextrose, isomalulose, polymaltose, arabinogalactan, palatinose, starches, starch hydrolysates, hydrogenated starch hydrolysates, partially hydrolyzed starch, dextrans, hydrogenated hexoses, fructooligosaccharides, sorbitol, xylitol, mannitol, maltitol, galactitol, isomalt, and mixtures thereof.

44. The process of sweetening a chewing gum composition according to claim 35, wherein said chewing gum composition further comprises an organic acid selected from the group consisting of citric acid, malic acid, tartaric acid, and mixtures thereof.

45. The process of sweetening a chewing gum composition according to claim 35, wherein said chewing gum composition further comprises a flavor selected from the group consisting of essential oils, synthetic flavors, oils derived from plants and fruits, citrus oils, fruit essences, peppermint oil, spearmint oil, clove oil, oil of wintergreen, anise, cinnamon, tutti frutti, strawberry, raspberry, lemon, orange, artificial flavorings, and combinations thereof.

46. A process of extending the sweetness of a chewing gum composition comprising the step of:

sweetening a chewing gum composition with N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester according to claim 28.

47. A process of extending the flavor of a chewing gum composition comprising the step of:

sweetening a chewing gum composition with N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester according to claim 28.

48. A process of reducing the amount of flavor ingredient required to produce a full-flavored chewing gum composition and to maintain a level of flavor intensity comparable to a full-flavored chewing gum composition comprising the step of:

sweetening a chewing gum composition with N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester according to claim 28.

49. The process of sweetening a chewing gum composition according to claim 28, wherein said chewing gum composition further comprises a rapid release sweetener.

50. The process of sweetening a chewing gum composition according to claim 49, wherein the rapid release sweetener has a sweetness intensity maximum within the first 2 minutes of chewing.

51. The process of sweetening a chewing gum composition according to claim 49, wherein the rapid release sweetener is selected from the group consisting

of sucrose, mannitol, fructose, high fructose corn syrup, sorbitol, dextrose, corn syrup solids, hydrogenated starch hydrolysates, invert sugar, fructose, xylitol, and combinations thereof.

52. The process of sweetening a chewing gum composition according to claim 28, wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester is encapsulated by or admixed with an agent.

53. The process of sweetening a chewing gum composition according to claim 52, wherein the agent is selected from the group consisting of cellulose, cellulose derivatives, hydroxypropylmethyl cellulose, stearic acid, shellac, polyethylene wax 500, zein, sterine 27, alginates, gelatin, starches, proteins, sugars, sugar alcohols, complex carbohydrates, gums, hydrocolloids, gellan gum, polydextrose, polywax, hydrogenated starch hydrolyzate, polyvinyl acetate, xanthan gum, carrageenan, dextrose, malic acid, maltodextrin, gum arabic and combinations thereof.

2025-12-17 14:17:00

ABSTRACT

5 The use of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester (neotame) as a sweetener in chewing gum compositions is disclosed. The chewing gums have been found to have a clean sweet taste profile with an extremely low level of sweetener usage. The use of neotame as a sweetener in chewing gum compositions also provides an extension of both
10 sweetness and flavor as compared to that expected from the use of other sweetener ingredients.

FIG. 1




FIG. 2




FIG. 3





FIG. 4



COMBINED DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION

(Page 1)

As a below named inventor, I hereby declare that

My residence, post office address and citizenship are as stated below next to my name

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

N - [N- (3, 3 - DIMETHYLBUTYL) - L- α -ASPARTYL] - L- PHENYLALANINE 1-METHYL
ESTER AS A SWEETENER IN CHEWING GUM

the specification of which ☒ is attached hereto ☐ was filed on _____ as United States
Application No. or PCT International Application No. _____
and was amended on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended
by any amendment referred to above

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR §1.56

I hereby claim foreign priority benefits under 35 U.S.C. §119(a)-(d) or §365(b), of any foreign application(s) for patent or inventor's
certificate, or §365(a) of any PCT international application which designates at least one country other than the United States, listed below
and have also identified below any foreign application for patent or inventor's certificate, or PCT international application having a filing date
before that of the application on which priority is claimed

Country	Application No.	Filed (Day/Mo./Yr.)	(Yes/No) Priority Claimed
---------	-----------------	---------------------	------------------------------

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s), or §365(c) of any PCT international application
designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the
prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty
to disclose information which is material to patentability as defined in 37 C.F.R. §1.56 which became available between the filing date of the
prior application and the national or PCT international filing date of this application.

Application No.	Filed (Day/Mo./Yr.)	Status (Patented, Pending, Abandoned)
-----------------	---------------------	---------------------------------------

I hereby claim the benefit under 35 U.S.C. §119(e) of United States Provisional Application No. 60/112,915, filed 18 December
1998

I hereby appoint the practitioners associated with the firm and Customer Number provided below to prosecute this application and
to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to the address
associated with that Customer Number

FITZPATRICK, CELLA, HARPER & SCINTO
Customer Number: 05514

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief
are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are
punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements
may jeopardize the validity of the application or any patent issued thereon

Full Name of Sole or First Inventor SUBBARAO V. PONAKALA

Inventor's signature _____

Date _____ Citizen/Subject of United States

Residence Elk Grove Village, Illinois

Post Office Address 981 Kentucky Lane, Elk Grove Village, IL 60007

COMBINED DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION

(Page 2)

Full Name of Second Joint Inventor JEANETTE ZIEGLER

Second Inventor's signature _____

Date _____ Citizen/Subject of United States

Residence Kenosha, Wisconsin

Post Office Address 1438 97th Avenue, Kenosha, WI 53144

Full Name of Third Joint Inventor, if any KENNETH PATTERSON

Third Inventor's signature _____

Date _____ Citizen/Subject of United States

Residence Lake Zurich, Illinois

Post Office Address 1264 William Drive, Lake Zurich, IL 60047

Full Name of Fourth Joint Inventor, if any STEVE A. SCHROEDER

Fourth Inventor's signature _____

Date _____ Citizen/Subject of United States

Residence Belvidere, Illinois

Post Office Address 6486 Hallen Avenue, Belvidere, IL 61008